

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

P^rMAR-PREGABALIN

Pregabalin Capsules,

Read this carefully before you start taking **MAR-PREGABALIN** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **MAR-PREGABALIN**.

Serious Warnings and Precautions

Taking MAR-PREGABALIN with opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.

What is MAR-PREGABALIN used for?

MAR-PREGABALIN is used in adults to treat pain caused by nerve damage due to:

- Diabetes
- Shingles

MAR-PREGABALIN is also used to treat pain associated with a condition called fibromyalgia (widespread pain).

How does MAR-PREGABALIN work?

MAR-PREGABALIN works by decreasing the number of nerve signals in your body. This helps to calm down oversensitive nerve cells, which helps to relieve pain.

What are the ingredients in MAR-PREGABALIN?

Medicinal ingredients: Pregabalin

Non-medicinal ingredients: The non-medicinal ingredients that make up the capsule core are pre-gelatinized starch and talc.

The capsule shells contain gelatin and titanium dioxide. In addition, the orange capsule shells contain iron oxide red.

The markings on the capsules are in black ink, which contains ammonia solution, black iron oxide, butyl alcohol, dehydrated alcohol, isopropyl alcohol, potassium hydroxide, propylene glycol, shellac, and water.

MAR-PREGABALIN comes in the following dosage forms:

Capsules: 25 mg, 50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 225 mg and 300 mg.

Do not use MAR-PREGABALIN if:

- You are allergic to pregabalin, or any other ingredient in MAR-PREGABALIN.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take MAR-PREGABALIN. Talk about any health conditions or problems you may have, including if you:

- have any kidney problems.
- have any breathing problems or respiratory disease.
- are pregnant, plan to become pregnant, or think you might be pregnant.
- are breastfeeding. MAR-PREGABALIN passes into breast milk and it is not known if it can harm your baby. You and your healthcare provider should discuss whether you should take MAR-PREGABALIN or breast-feed, but you should not do both.
- have ever had an allergic reaction to any other drug.
- have a history of heart disease or heart failure.
- have a history of lower gastrointestinal problems (e.g., constipation, blocked or paralysed bowel), or you are taking medication(s) that may cause constipation.
- have a history of addiction or substance abuse.

Other warnings you should know about:

Pregnancy: MAR-PREGABALIN should not be taken during pregnancy. If you become pregnant while taking MAR-PREGABALIN, tell your healthcare professional **right away**. If you are taking MAR-PREGABALIN and are of childbearing age, you must use an effective method of birth control. If you take MAR-PREGABALIN during your first trimester of pregnancy, it can cause major birth defects in your unborn child.

Pregnancy Registry: If you become pregnant while taking MAR-PREGABALIN, talk to your healthcare professional about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334. The purpose of this registry is to collect information about the safety of anticonvulsant medicines during pregnancy. Information on the registry can also be found at the website <http://www.aedpregnancyregistry.org/>.

Eye Problems: If you experience any changes in your vision while taking MAR-PREGABALIN, tell your healthcare professional right away. If you already have eye problems, your healthcare professional may recommend that you have more frequent eye exams.

Stopping Your Medication: Do NOT stop taking MAR-PREGABALIN without discussing it with your healthcare professional first. Stopping your treatment abruptly may cause you to experience unwanted side effects. If you have any concerns with your treatment, talk to your healthcare professional.

Driving and Using Machines: MAR-PREGABALIN may cause you to feel dizzy or sleepy. Wait until you know how MAR-PREGABALIN affects you before performing tasks that require special attention.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with MAR-PREGABALIN:

- Alcohol

- Opioid medicines
- Benzodiazepines
- Medications used to treat diabetes, such as rosiglitazone, pioglitazone

How to take MAR-PREGABALIN:

- Take MAR-PREGABALIN exactly as your healthcare professional has told you to.
- Do NOT stop taking MAR-PREGABALIN abruptly or you may experience unwanted side effects. If you have concerns with your treatment, talk to your healthcare professional first.
- You can take MAR-PREGABALIN with or without food.

Usual dose:

Your healthcare professional has decided the best dose for you and will tell you how much to take and when to take it.

Overdose:

If you think you, or a person you are caring for, have taken too much MAR-PREGABALIN, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose by a few hours, take it as soon as you remember. If it is close to your next dose, do NOT take the missed capsules. Simply wait and restart MAR-PREGABALIN with your next scheduled dose.

What are possible side effects from using MAR-PREGABALIN?

These are not all the possible side effects you may have when taking MAR-PREGABALIN. If you experience any side effects not listed here, tell your healthcare professional.

Side effects may include:

- Dizziness
- Sleepiness
- Headache
- Nausea
- Dry mouth
- Increased cough
- Weight gain
- Trouble concentrating
- Forgetfulness
- Lack of energy
- Muscle weakness
- Constipation; talk with your healthcare professional about ways to prevent constipation when you start using MAR-PREGABALIN

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
COMMON			
Blurred vision		✓	
UNCOMMON			
Edema: unusual swelling of the arms, hands, legs, feet and ankles, face or airway passages		✓	
Angiodema: swelling of the face, mouth, lips, gums, neck or throat, trouble breathing, diarrhea, nausea or vomiting			✓
Allergic Reaction: difficulty swallowing or breathing, wheezing, feeling sick to your stomach and throwing up, hives or rash, swelling of the face, lips, tongue or throat			✓
Decrease in the amount of urine		✓	
Suicidal thoughts or actions		✓	
RARE			
Kidney Failure: confusion, itchiness or rashes, puffiness in your face and hands, swelling in your feet or ankles, decreased amount of urine, weight gain			
VERY RARE			
Seizures (fits): uncontrollable shaking with or without loss of consciousness.			✓
Severe Skin Reactions: fever, severe rash, swollen lymph glands, flu-like feeling, blisters and peeling skin that may start in and around the mouth, nose, eyes and genitals and spread to other areas of the body, yellow skin or eyes, shortness of breath, dry cough, chest pain or discomfort, feeling thirsty, urinating less often, less urine			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Store at room temperature (15°C to 30°C) in the package it came in.

Keep out of reach and sight of children.

If you want more information about MAR-PREGABALIN:

- Talk to your healthcare professional
 - Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website www.marcanpharma.com, or by calling 1-855-627-2261.

This leaflet was prepared by Marcan Pharmaceuticals Inc.

Last Revised: September 13, 2023