

**Table 3:** Table showing total skin lesion referrals grouped into malignant, non-malignant or inflammatory categories and their respective commonest (greater than 1%) dermatologist clinical diagnosis.

	<b>Dermatologist clinical diagnosis</b>	<b>Percentage (total 233)</b>
<b>Malignant skin lesions 30%</b>	Basal cell carcinoma (BCC)	12%
	Melanoma	9%
	Squamous cell carcinoma (SCC)	9%
<b>Malignant precursor 14%</b>	Actinic keratosis	14%
<b>Benign lesions 23%</b>	Seborrheic keratosis	9%
	Naevi	9%
	Wart	4%
	Molluscum contagiosum	1%
<b>Inflammatory lesions 33%</b>	Eczema	1%
	Acne	1%
	Granuloma annulare	1%
	Prurigo	1%
	Tinea	1%
	Other (each <1%)	28%

**Table 4:** Comparison of commonest benign, premalignant and malignant keratinocyte and pigmented lesion diagnoses between a general practitioner with a special interest in skin cancer (GPwSISC) and other practitioners.

<b>Clinical diagnosis</b>	<b>General practitioner with a special interest in skin cancer (GPwSISC)</b>	<b>Percentage</b>	<b>Other practitioner</b>	<b>Percentage</b>
Superficial basal cell carcinoma (BCC)	2	1%	2	2%
Basal cell carcinoma (BCC)	19	14%	8	9%
Squamous cell carcinoma in situ	0	0%	4	4%
Squamous cell carcinoma (SCC)	8	6%	3	3%
Melanoma	13	9%	2	2%
Actinic keratosis	18	13%	21	23%
Other suspicious malignancy (eg, Merkel's, lymphoma, amelanotic melanoma)	6	4%	2	2%
Naevi	12	9%	6	7%
Lentigo	7	5%	1	1%
Seborrheic keratosis	3	2%	18	20%
Other benign lesions	53	37%	25	27%
<b>Total lesions</b>	<b>141</b>	<b>100%</b>	<b>92</b>	<b>100%</b>