

Cabotegravir/Rilpivirine (Cabenuva)



Provider Order Form rev. 10/30/2023

PATIENT INFORMATION

Referral Status: New Referral Updated Order Order Renewal

Patient Name: _____ DOB: _____ Patient Phone: _____

Patient Address: _____ Patient Email: _____

Allergies: _____ NKDA Weight (lbs/kg): _____ Height (in/cm): _____

Sex: M / F Date of Last Infusion: _____ Next Due Date: _____ Preferred Location: _____

DIAGNOSIS *(Please provide ICD-10 code in space provided)*

HIV infection (to replace current stable antiretroviral regimen): _____

Other: _____ Description: _____

THERAPY ADMINISTRATION & DOSING

- Induction: Administer Cabotegravir 600mg /Rilpivirine 900mg IM as separate gluteal injections (on opposite sides at least 2cm apart) on week 0.
- Maintenance: administer Cabotegravir _____ mg /Rilpivirine _____ mg IM as separate gluteal injections (on opposite sides at least 2cm apart) one month after initial injection followed by every _____ months.
- Monitor patient for 10 minutes after every injection. Vital signs prior to discharge.

NURSING

- Hold infusion and notify provider if patient has not adhered to the monthly or every other monthly regimen.
- Provide nursing care per Nursing Procedure, including Hypersensitivity Reaction Management Protocol and post-procedure observation.

PRE-MEDICATION ORDERS

Other: _____

ADDITIONAL ORDERS

PROVIDER INFORMATION

Preferred Contact Name: _____ Preferred Contact Email: _____

Ordering Provider: _____ Provider NPI: _____

Referring Practice Name: _____ Phone: _____ Fax: _____

Practice Address: _____ City: _____ State: _____ Zip Code: _____

REQUIRED DOCUMENTATION CHECKLIST *(Additional documentation required for processing and insurance approval)*

Required Documentation: Patient demos, copy of front and back of primary and secondary insurance, 2 most recent OVN including treatment failures or contraindications, Confirmation of HIV-1 diagnosis/Greater than 18 years of age, Virologic suppression for at least 6months (HIV-1 RNA<50 copies/ml), Documentation of adherence to oral therapy, No history or TF/resistance to medication, Complete medication profile (contraindication of use with cabamazepime, oxycarbazepine, phenobarbital, phenytoin, Rifabutin, Rifampin, Rifapentine, steroids, St. John's Wort)

Required Labs: LFT, Renal function

Provider Name *(print)*

Provider Signature

Date

Order valid for one year unless otherwise indicated. IV solutions/diluents may be substituted as allowed per manufacturer's instructions as necessitated by product availability.