

Ravulizumab-cwvz (Ultomiris)

Provider Order Form rev. 10/30/2023

PATIENT INFORMATION

Referral Status: New Referral Updated Order Order Renewal

Patient Name: _____ DOB: _____ Patient Phone: _____

Patient Address: _____ Patient Email: _____

Allergies: _____ NKDA Weight (lbs/kg): _____ Height (in/cm): _____

Sex: M / F Date of Last Infusion: _____ Next Due Date: _____ Preferred Location: _____

DIAGNOSIS (Please provide ICD-10 code in space provided)

Myasthenia Gravis (anti-acetylcholine receptor antibody positive): _____

Other: _____ Description: _____

REQUIRED INFORMATION

Must have meningococcal vaccine at least 2 weeks prior to infusions. If no vaccine, referring provider must provide 2 weeks of antibacterial drug prophylaxis

THERAPY ADMINISTRATION & DOSING

Administer Ultomiris IV over 1 hour (**Choose one**):

Weight 40-60kg:

- Loading: 2400mg (in 24ml NS) at week 0, followed by 3000mg (in 30ml NS) at week 2
- Maintenance: 3000mg (in 30ml NS) every 8 weeks

Weight 60-100kg:

- Loading: 2700mg (in 27ml NS) at week 0, followed by 3300mg (in 33ml NS) at week 2
- Maintenance: 3300mg (in 33ml NS) every 8 weeks

Weight 100kg or more:

- Loading: 3000mg (in 30ml NS) at week 0, followed by 3600mg (in 36ml NS) at week 2
- Maintenance: 3600mg (in 36ml NS) every 8 weeks

Switching from Eculizumab: Administer loading dose 2 weeks after last dose of eculizumab followed by maintenance dose every 8 weeks

Monitor Patient for 60mins after every infusion

DO NOT mix 300mg/30ml vials with other concentrations. OK to mix 300mg/3ml and 1100mg/11ml together if needed.

LABORATORY ORDERS

Other: _____

PRE-MEDICATION ORDERS

- Tylenol 500mg / 650mg PO
 Loratadine 10mg PO
 Pepcid 20mg PO / IVP
 Benadryl 25mg / 50mg PO / IVP
 Solumedrol 40mg / 125mg IVP
 Other: _____

NURSING

- Hold infusion and notify provider for:
- abnormal vital signs or signs/symptoms of infection or Meningitis
 - New or worsening headache or altered mental status
- Record vitals before infusion then every 30mins until patient discharges. If reactions occur, slow or stop infusion
- Provide nursing care per Nursing Procedure, including Hypersensitivity Reaction Management Protocol and post-procedure observation.

ADDITIONAL ORDERS

PROVIDER INFORMATION

Preferred Contact Name: _____ Preferred Contact Email: _____

Ordering Provider: _____ Provider NPI: _____

Referring Practice Name: _____ Phone: _____ Fax: _____

Practice Address: _____ City: _____ State: _____ Zip Code: _____

REQUIRED DOCUMENTATION CHECKLIST (Additional documentation required for processing and insurance approval)

Required Documentation: Patient demos, copy of front and back of primary and secondary insurance, 2 most recent OVN including treatment failures or contraindications, EMG results, MRI results

Required Labs: AChR antibody, MuSK antibodies, CRP, ESR

Provider Name (print)

Provider Signature

Date