## Iron Infusion (Feraheme, Venofer, Monoferric, Injectafer)

Provider Order Form rev. 12/19/2023



PATIENT INFO	RMATION	<b>Referral Status:</b>	🗆 New R	eferral	Updated Orde	er 🛛 Order Renewal	
Patient Name:			DOB:		Patient Pho	ne:	
Patient Address:		Patient Email:					
Allergies:			□ NKDA	Weigh	t (lbs/kg):	Height (in/cm):	
Sex: 🗆 M / 🗆 F	Date of Last Infusion:	Next Due Date	2:	Pref	erred Location:		
DIAGNOSIS (P	lease provide ICD-10 co	de in space provided)					
Iron deficiency A	nemia:	chronic kidney disease:		CKD sta	ge required:		
Other:		Description:					
THERAPY ADMINISTRATION (Choose one)			LABORATORY ORDERS				
List in order of pre	luct as required by patient' ference:,, uct only (subject to prior a	uthorization)	<ul> <li>At least one month post last infusion of iron, draw CBC with diff, ferritin, Iron, saturation, TIBC.</li> <li>Phosphorus (<i>indicated with injectafer</i>)</li> <li>Other:</li></ul>				
<ul> <li>✓ Monitor patient infusion.</li> <li>DOSING &amp; FR Venofer Dose: □ 1 200mg in 100ml N and infuse over 90</li> <li>Venofer Frequence □ every □ every</li> </ul>	PRE-MEDICATION ORDERS  Tylenol 500mg / 650mg PO Loratadine 10mg PO Pepcid 20mg PO / IVP Benadryl 25mg / 50mg PO / IVP Solumedrol 40mg / 125mg IVP Other:						
<ul> <li>Feraheme Dose &amp; Frequency:</li> <li>□ administer 510mg IV x2 doses (Separated by 3-8 days). Mix in 100ml NS and infuse over 15-30 minutes.</li> <li>Injectafer Dose &amp; Frequency:</li> <li>□ Pts over 50kg, administer 750mg IV on day 0 and day 7</li> <li>□ Pts under 50kg, administer 15mg/kg IV = mg on day 0</li> </ul>			<ul> <li>Hold infusion and notify provider for history of allergy to IV iron</li> <li>Place patient in reclined or semi-reclined position.</li> <li>Use with caution in patients with hypotension (<i>feraheme/venofer</i>)</li> <li>Provide nursing care per Nursing Procedure, including</li> <li>Hypersensitivity Reaction Management Protocol and post-procedure observation</li> </ul>				

## **ADDITIONAL ORDERS**

## **PROVIDER INFORMATION**

**Monoferric Dose & Frequency** 

single dose. Dilute in 100ml NS

Mix in 250ml NS and infuse over 30 minutes.

□ Pts under 50kg, administer 20mg/kg IV =

□ Pts over 50kg, administer 1000mg IV over at least 20min as

least 20mins as single dose. Dilute to final concentration of

and day 7

1mg/ml

Preferred Contact Name:	Pref	Preferred Contact Email:			
Ordering Provider: Provider NPI:					
Referring Practice Name:	Phone:	Fax:			
Practice Address:	City:	State:	Zip Code:		

## REQUIRED DOCUMENTATION CHECKLIST (Additional documentation required for processing and insurance approval)

**Required Documentation:** Patient demos, copy of front and back of primary and secondary insurance, 2 most recent OVN including treatment failures or contraindications with oral iron, Reason for anemia (by indication)

Required Labs: Kidney function, CBC, Ferritin, Iron, TIBC, Iron saturation, Iron within the last 4 weeks.

mg over at

Provider Name (print)

**Provider Signature** 

Date