
Ketamidor[®]

Ketamidor 100 mg/ml solution for injection



Ketamidor is indicated to be used to induce anaesthesia in the following circumstances:

- · In combination with detomidine in the horse.
- In combination with xylazine in the horse, in cattle, dog and in the cat.
- · In combination with azaperone in the pig.
- In combination with medetomidine in the dog and cat.
- · In combination with diazepam in the dog.

Ketamidor is indicated for use as a sole agent for restraint and minor surgical procedures in the cat, where muscle relaxation is not required.

Recovery usually is complete after 2 hours, but may occasionally take longer.

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.

Shelf life after first opening the immediate packaging: 28 days.



Active	Ketamine (as hydrochloride)	
Species	Horse, Cattle, Pig, Dog, Cat	
Routes of Administration	IV - Horse, Cattle, Dog, Cat IM - Pig, Dog, Cat SC - Cat	
Pack Size	10ml, 50ml	
Legal Category	VPO, POM-V	



Ketamidor[®] Mono Dosage Guide



Substances & Combinations	Aim	Dosages in mg/kg			Route
	Λ	K	М	X	
Ketamine + Detomidine	Anaesthesia - give alpha- 2 agonist first followed by ketamine fast IV	2.2	0.02	-	IV IM
Ketamine + Xylazine		2.2	-	1.1	IV IM

Pre-medication with a sedative is required for a sufficient anaesthetic effect, ensure adequate time for sedative to take effect before induction.



Substances & Combinations	Aim	Dosages in mg/kg		Route
	Angesthesia	K	X	
Ketamine + Xylazine*	give xylazine first followed by ketamine	2	0.14	IV
		5	0.22	IM

^{*} The lower end of the stated dose range should be used when administering xylazine via the intravenous route

To avoid uncontrolled lying down and possible symptoms of excitation or for potentiation of anaesthesia a sedative premedication is recommended. To avoid hypoxia due to lateral or dorsal recumbency, oxygen can be administered through a nasal tube.



Substances & Combinations	Aim	Dosages in mg/kg		Route
		K	X	
Ketamine +		15	2	IV.
Azaperone		20	2	IV

Doses based on those in product SPC. Doses should be tailored to individual patient depending on clinical status, indication, combination of drugs used and required level of sedation.

The dosage amounts are based on the dose rate in the product SPC. Dosage amounts should be tailored to the individual patient depending on the status, indication, combination of drugs used and the required level of sedation. Ketamidor 100 mg/ml solution for injection. Contains: Ketamine. Vm 22080/4004.

Refer to product packaging and leaflets for full indications, side effects, precautions, warnings and contra-indications.

Use medicines responsibly.

Further information can be found on the datasheet, SPC or at www.chanellepharma.com. Distributed by: Chanelle Animal Health, Loughrea, Co. Galway, Ireland. Copyright © Chanelle 2020. All rights reserved.