



Kauri

Agathis australis

INTRODUCTION AND METHODS

Reasons for planting native trees include the enhancement of plant and animal biodiversity for conservation, establishment of a native cover on erosion-prone sites, improvement of water quality by revegetation of riparian areas and management for production of high quality timber. Significant areas of the New Zealand landscape, both urban and rural, are being re-vegetated using native species. Many such plantings are on open sites where the aim is to quickly achieve canopy closure and often includes the planting of a mixture of shrubs and tree species concurrently. Previously, data have been presented showing the potential above- and below-ground growth performance of eleven native plant species considered typical early colonisers of bare ground, particularly in riparian areas (<http://icm.landcareresearch.co.nz/research/land/Trial1results.asp>). In this current series of posters we present data on the growth performance of six native conifer (kauri, rimu, totara, matai, miro, kahikatea) and two broadleaved hardwood (puriri, titoki) species most likely to succeed the early colonising species to become a major component in mature stands of indigenous forest (<http://icm.landcareresearch.co.nz/research/land/Trial2.asp>). Data on the potential above- and below-ground early growth performance of colonising shrubby species together with that of conifer and broadleaved species will help land managers and community groups involved in re-vegetation projects in deciding the plant spacing and species mix most appropriate for the scale of planting and best suited to site conditions.

Data are from a trial established in 2006 to assess the relative growth performance of native conifer and broadleaved hardwood tree species. Ten plants were extracted each year for 5 years following establishment and their above- and below-ground growth parameters measured.




Side view of 5-year old root system
(see text box for dimensions)




View of canopy and root system of a 5-year old plant (see text box for dimensions)

RESULTS

TREE HEIGHT



CANOPY AND ROOT SPREAD



DISTRIBUTION AND SITE PREFERENCES


Occurrence	North Cape, westward to Tauranga and southward to Kawhia
Local occurrence	Lowland and montane forests, sea level to 360 m (800 m on Mt Moehau)
Preferred soils	Good growth rates on fertile colluvial and weathered volcanic soils, tolerate infertile and ultra-infertile soils
Moisture	Tolerant of periodic dry conditions hence survives on dry, skeletal ridges, intolerant of persistently wet sites
Properties	Requires warm temperatures and relatively high light conditions. Resistant to drought and lack of nutrients, long lived

SUMMARY OF GROWTH CHARACTERISTICS AT AGE 5


Mean Height	1.55 m
Mean canopy	0.80 m
Mean root spread	0.52 m
Mean max. root depth	0.36 m
Mean above-ground biomass	0.25 kg
Mean below-ground biomass	0.07 kg
Root:shoot ratio	0.27

Notes: Able to occupy a site for several generations. Best growth on lower slopes where trees have survived competition from broadleaf species. Makes effective use of nutrients and water and is thus suitable for poorer soil types. Planted Kauri seems to grow better on moist, fertile soils than naturally regenerated Kauri. Benefits from weed control in early years following establishment. Grows up to 30 m tall.


ROOT DEPTH




ROOT LENGTH DISTRIBUTION



BIOMASS



TOTAL PLANT BIOMASS



ROOT BIOMASS DISTRIBUTION

