



**Read
Before
Use**

As a precaution, gloves and eye protection should be worn when handling substances and using test units. Do not puncture any part of unit. Do not remove probe once a capsule is broken. In the unlikely event of a spill, rinse affected area thoroughly with water. Keep out of reach of children. All results are presumptive.

Test Preparation	1	Remove the Smart-Tip™ probe from test unit.	2		Sample suspected substance or surface.	
			Substance & Trace Material		Liquid	
			Swipe the paper of the Smart-Tip™ probe across the suspect substance or surface.		Place the absorbent paper of Smart-Tip™ probe into liquid. Make sure paper is wet.	

Test Preparation	3	Lock Smart-Tip™ probe into test unit. Press firmly.	STOP Do not continue unless probe is locked below the Flex Pins and cannot be removed from the test unit.	Identification	4	With thumb and forefinger , squeeze chamber 1 until capsule breaks.	5	Gently shake test unit for 2-3 seconds.

Identification	6	With thumb and forefinger , squeeze chamber 2 until capsule breaks.	7	Gently shake test unit for 2-3 seconds.

Results	 	The appearance of an orange or yellow color in either the front or back window indicates the presence of Peroxide .
		If color develops, proceed to Disposal. If no color develops, proceed to Step 8.

Chamber 3 Identification	8	With thumb and forefinger , squeeze chamber 3 until capsule breaks.	9	Gently shake test unit for 10 seconds.

Chamber 3 Results	 	The appearance of an orange or yellow color in either the front or back window indicates the presence of TATP or HMTD .
		If no orange or yellow color develops, the above listed substances are <u>not</u> present.



Disposal

The test kit is safe to dispose of in the trash.

0739