

General Screening
Drugs
Test Kits





Read Before Use As a precaution, gloves and eye protection should be worn when handling substances and using test units. Do not puncture any part of unit. Do not remove probe after breaking the ampule. In event of a spill, rinse affected area thoroughly with water. Keep out of reach of children. Store the tubes in a dark place, not exposed to light. All results are presumptive. If necessary, further testing by a competent drug testing lab can provide additional identification.

est Preparation

Remove the SWIPPER<sup>TM</sup> Cap from the test unit.

Ensure that the test tube remains upright in order to avoid the ampule falling out of the tube.

Sample the suspected substance with the SWIPPER<sup>TM</sup> Cap.



**Test Preparation** 

Return the SWIPPERTM Cap to the test tube. Press firmly.



## **STOP**

Do not continue unless the SWIPPER<sup>TM</sup> Cap is properly secured.

Identification

With thumb and forefinger, squeeze the test tube until the ampule breaks.



Gently shake test up and down to ensure that the liquid touches the Cap.



Results

The immediate appearance of a purple-red color indicates the presence of one of the following drugs:

Heroin, Amphetamine, Methamphetamine, Ecstasy/MDMA, Benzedrone, Buphedrone, Carfentanyl, Cathinone, DMT, EAPB, Fentanyl, Ketamine, LSD, mCPP, MDMAI, MDPBP, MDPV, Mephedrone, Methadone, Methcathinone, Methylone, Mexedrone, MPA, PCP, Pentedrone, Pentylone, PMA, PMMA, Psilocybin, 2-aminoindane (2-AI), other synthetic/designer drugs.



The immediate appearance of a green or blue-green color indicates the presence of **Cocaine** or **Crack**.



If none of the above colors develop, the above listed drugs are not present.



The test kit is safe to dispose of in the trash.

Donto



Manufactured by IDenta Ltd www.identa-corp.com Patented Technology • Made in Israel LOT:

EXP: