

(Peroxide-TATP Test) Peroxide




As a precaution, gloves and eye protection should be worn when handling substances and using test units. If a spill occurs, thoroughly rinse affected area with water. Keep out of reach of children. Store the tubes in a dark place, not exposed to light. All results are presumptive.

This test consists of two test tubes, which much be applied in order. Test in sequence-Tube #1—Tube #2. Since the tubes contains enough liquid to test many samples, do not dispose of the tube until all desired samples are tested. For additional tests with the same tubes, take a new sheet of test paper from the pad to test the next sample.

Remove a sheet of test paper for every test from the dispenser.













Initial Use: With thumb and forefinger, squeeze Tube #1 until the ampule breaks.



Open the cap and apply a drop to the sampled area of the test paper.






The appearance of an orange or yellow color indicates the presence of Peroxide.



If an orange or yellow color appears, testing is completed. If no color develops immediately, proceed to the next step.





Open the cap and apply a drop to the same place as Step





The appearance of an orange or yellow color indicates the presence of TATP (Triacetone Triperoxide) or HMTD (Hexamethylene Triperoxide Diamine).



If no orange or yellow color develops, the above substances are not present.



The tubes contain enough liquid for multiple tests. Replace cap after use for storage and further use. When finished, the tubes can be discarded in the trash.

IDenta

World Leader in Forensic Products

Manufactured by IDenta Ltd www.identa-corp.com Patented Technology • Made in Israel LOT: EXP: